

Combinatorial properties of the numbers of tableaux of bounded height

Marilena Barnabei, Flavio Bonetti, and Matteo Silimbani *

Abstract

We introduce an infinite family of lower triangular matrices $\Gamma^{(s)}$, where $\gamma_{n,i}^s$ counts the standard Young tableaux on n cells and with at most s columns on a suitable subset of shapes. We show that the entries of these matrices satisfy a three-term row recurrence and we deduce recursive and asymptotic properties for the total number $\tau_s(n)$ of tableaux on n cells and with at most s columns.

1 Introduction

The first simple expressions of the number of standard Young tableaux of given shape were given by Frobenius and Young (see [6] and [15]) and by Frame-Robinson-Thrall [5]. More recently, the number of standard Young tableaux has been studied according to the height of their shape. Regev [12] gave asymptotic values for the numbers $\tau_s(n)$ of standard Young tableaux whose shape consists of n cells and at most s columns and Stanley (see [14]) discussed the algebraic or differentiably finite nature of the corresponding generating functions. More recently, many authors entered this vein. For example, D.Gouyou-Beauchamps gave both exact forms in [13] and recurrence formulas (see [9]) for the numbers $\tau_s(n)$ when $s \leq 5$ by combinatorial tools, while F.Bergeron and F.Gascon [4] found some recurrence formulas for the numbers $\tau_s(n)$ by analytic argumentations.

In this paper we present a different approach to the subject. Our starting point is the remark that the elements of the Ballot matrix introduced

*Department of Mathematics - University of Bologna

by M.Aigner in [1] correspond bijectively to the integers counting standard Young tableaux of a given shape with at most 2 columns. Firstly, we arrange the entries of the Ballot Matrix in a new lower triangular matrix A in such a way that the entries of the n -th row count standard Young tableaux with precisely n cells. The integer $\tau_2(n)$ can be therefore recovered by summing the elements of the n -th row of A .

In the following sections, we extend the results to the general case of standard Young tableaux with at most s columns. For every fixed $s \in \mathbb{N}$ we define an infinite matrix $\Gamma^{(s)}$ whose (n, i) -th entry is the total number of standard Young tableaux with at most s columns and such that the difference between the length of the second and of the third column is i . The total number $\tau_s(n)$ of tableaux on n cells with at most s columns is the n -th row sum of the matrix $\Gamma^{(s)}$. The matrix $\Gamma^{(s)}$ presents a three-term row-recurrence property, which yields a recurrence law satisfied by the integers $\tau_s(n)$. This recurrence allows to get a combinatorial interpretation of the asymptotic behaviour of the ratio

$$\frac{\tau_s(n)}{\tau_s(n-1)}.$$

In Section 3 we treat more extensively the case $s = 3$, in order to exhibit recursive and asymptotic properties for the integers $\tau_3(n) = M_n$, where M_n denotes the n -th Motzkin number.

2 The two-column case

In the following, we will denote the shape of a standard Young tableau T as a list containing the length of the first, second, ..., last column of T .

We first consider the standard Young tableaux whose shape consists of at most two columns. We define an infinite, lower triangular matrix $A = (\alpha_{n,i})$, where $\alpha_{n,i}$ is the number of standard Young tableaux whose shape corresponds to the list $(n-i, i)$, with $i \in \mathbb{N}$.

The matrix A can be recursively constructed as follows:

Proposition 1 *The matrix A is defined by the following initial conditions:*

- i) $\alpha_{i,0} = 1$ for every $i \geq 0$;
- ii) $\alpha_{1,i} = 0$ for every $i > 0$;

iii) $\alpha_{n,i} = 0$ for every $i > \lfloor \frac{n}{2} \rfloor$

and by the recurrence:

$$\alpha_{n,i} = \alpha_{n-1,i} + \alpha_{n-1,i-1} \quad \text{for } n \geq 1, 1 \leq i \leq \lfloor \frac{n}{2} \rfloor. \quad (1)$$

Proof The initial conditions follow immediately by the definition of the matrix A . In order to prove the recurrence, remark that a Young tableau with n cells can be obtained from a tableau with $n-1$ cells by adding a new box containing the symbol n in a corner position. In particular, in a Young tableau of shape $(n-i, i)$, $i > 0$, the symbol n can be placed in at most two corner cells. Hence, the entry $\alpha_{n,i}$ is the sum of the two integers $\alpha_{n-1,i-1}$ and $\alpha_{n-1,i}$, that count the Young tableaux of appropriate shape. $\square$

The recurrence formula (1) has as an immediate fallout the following result:

Corollary 2 *The entries of the matrix A satisfy the following columnwise recurrence:*

$$\alpha_{n,i} = \sum_{h=2i-1}^{n-1} \alpha_{h,i-1}$$

for every $n, i > 0$.

We remark that the elements of the matrix A correspond bijectively to the entries of the Ballot Matrix defined by Aigner in [1]. More precisely, the Ballot Matrix $\tilde{A} = (\tilde{a}_{i,j})$ can be obtained rearranging the entries of the matrix A as follows:

$$\tilde{a}_{j,k} = \alpha_{2j-k,j-k}.$$

As a consequence, for every n , the number of standard Young tableaux with two columns of the same length n is the n -th Catalan number C_n :

$$\alpha_{2n,n} = C_n \quad (2)$$

.

Denote by $\tau_2(n)$ the number of Young tableaux with n cells and at most two columns. Obviously, we have:

$$\tau_2(n) = \sum_{i=0}^{\lfloor \frac{n}{2} \rfloor} \alpha_{n,i}.$$

In the next theorem we deduce a recurrence formula for the sequence $\tau_2(n)_{n \geq 1}$:

Theorem 3 *The integers $\tau_2(n)$, $n \geq 1$, satisfy the recurrence*

$$\tau_2(n) = 2\tau_2(n-1) - E(n-1) \cdot C_{\frac{n-1}{2}}, \quad (3)$$

where C_i denotes the i -th Catalan number and

$$E(n) = \begin{cases} 1 & \text{if } n \text{ is even} \\ 0 & \text{otherwise.} \end{cases}$$

Proof By Proposition 1, we have:

$$\begin{aligned} \tau_2(n) &= \sum_{i=0}^{\lfloor \frac{n}{2} \rfloor} a_{n,i} = \sum_{i=0}^{\lfloor \frac{n}{2} \rfloor} a_{n-1,i-1} + \sum_{i=0}^{\lfloor \frac{n}{2} \rfloor} a_{n-1,i} = \\ &= 2\tau_2(n-1) - E(n-1) \cdot \alpha_{n-1, \frac{n-1}{2}} = 2\tau_2(n-1) - E(n-1) \cdot C_{\frac{n-1}{2}}, \end{aligned}$$

as desired. □




Figure 1: The recurrence formula for the entry $\alpha_{8,3}$.

The preceding result yields an alternative proof of the following theorem, originally proved by Regev in [12]:

Theorem 4 *The number of standard Young tableaux with exactly n cells and at most two columns is*

$$\tau_2(n) = \binom{n}{\lfloor \frac{n}{2} \rfloor}.$$

Proof It is well known that the central binomial coefficients $\binom{n}{\lfloor \frac{n}{2} \rfloor}$ satisfy the recurrence (3). Remarking that $\tau_2(0) = 1 = \binom{0}{0}$, we have the assertion. $\square$

Moreover, identity (3) yields the following asymptotic property of the integers $\tau_2(n)$:

Proposition 5 *We have:*

$$\lim_{n \rightarrow \infty} \frac{\tau_2(n)}{\tau_2(n-1)} = 2$$

Proof By (3) we get:

$$\frac{\tau_2(n)}{\tau_2(n-1)} = \frac{2\tau_2(n-1) - E(n+1) \cdot C_{\frac{n-1}{2}}}{\tau_2(n-1)}.$$

This implies that the assertion is proved as soon as we show that

$$\lim_{n \rightarrow \infty} \frac{E(n+1) \cdot C_{\frac{n-1}{2}}}{\tau_2(n-1)} = 0.$$

If n is even, this identity trivially holds. If n is odd, the described correction term is negligible with respect to $\tau_2(n-1)$ as n goes to infinity. In fact, we get:

$$\lim_{n \rightarrow \infty} \frac{\frac{1}{n} \binom{n-1}{\frac{n-1}{2}}}{\binom{n-1}{\frac{n-1}{2}}} = \lim_{n \rightarrow \infty} \frac{1}{n} = 0,$$

as required. $\square$

3 The three-column case

We now extend the results of Section 2 to the case of standard Young tableaux with at most three columns. Define a matrix $B = (\beta_{n,i})$ such that the (n, i) -th entry of B is the total number of standard Young tableaux of shape $(n-i-2k, i+k, k)$, for every possible value of k . In other terms, the integer $\beta_{n,i}$ is the cardinality of the set $Y_{n,i}^3$ containing all standard tableaux on n

cells with at most 3 columns such that the difference between the second and third column is i .

The entries of the matrix B satisfy a three-term recurrence:

Proposition 6 *The integers $\beta_{n,0}$ satisfy the identity:*

$$\beta_{n,0} = \beta_{n-1,0} + \beta_{n-1,1}. \quad (4)$$

Moreover, for every $1 \leq i \leq \lfloor \frac{n}{2} \rfloor$, we have:

$$\beta_{n,i} = \beta_{n-1,i-1} + \beta_{n-1,i} + \beta_{n-1,i+1} - r_{n,i}, \quad (5)$$

where

$$r_{n,i} = \begin{cases} \text{number of tableaux of shape } (\frac{n+i-2}{3}, \frac{n+i-2}{3}, \frac{n-2i+1}{3}) & \text{if } n-2i \equiv_3 2 \\ 0 & \text{otherwise.} \end{cases}$$

Proof Let T be a standard Young tableau T with 3 columns of shape $(n-i-2k, i+k, k)$, for some k , and let T' be the tableau obtained from T by removing the cell containing the symbol n . Obviously, the tableau T' has either shape $(n-i-2k-1, i+k, k)$ or $(n-i-2k, i+k-1, k)$ or $(n-i-2k, i+k, k-1)$. This implies that the correspondence $T \mapsto T'$ maps the set $Y_{n,i}^3$ into the union $Y_{n-1,i-1}^3 \cup Y_{n-1,i}^3 \cup Y_{n-1,i+1}^3$. However, such map is not in general surjective. In fact, it is easy to check that the difference between the set $Y_{n-1,i-1}^3 \cup Y_{n-1,i}^3 \cup Y_{n-1,i+1}^3$ and the image of the map consists exactly of the tableaux of shape $(\frac{n+i-2}{3}, \frac{n+i-2}{3}, \frac{n-2i+1}{3})$. This proves (5). The first recurrence can be proved by the same arguments, observing that, in this case, the map defined above is surjective. $\square$

The preceding result yields a recurrence formula satisfied by the total number $\tau_3(n)$ of standard Young tableaux with n cells and at most 3 columns:

Theorem 7 *The numbers $\tau_3(n)$, $n \geq 3$, satisfy the recurrence*

$$\tau_3(n) = 3\tau_3(n-1) - R(n) - E(n-1) \cdot C_{\frac{n-1}{2}} - \beta_{n-1,0}, \quad (6)$$

where

$$R(n) = \sum_{i=0}^{\lfloor \frac{n}{2} \rfloor} r_{n,i}.$$

Proof By definition of $\beta_{n,i}$ we have:

$$\tau_3(n) = \sum_{i=0}^{\lfloor \frac{n}{2} \rfloor} \beta_{n,i}.$$

Hence,

$$\tau_3(n) = \sum_{i=0}^{\lfloor \frac{n}{2} \rfloor} \beta_{n-1,i-1} + \sum_{i=0}^{\lfloor \frac{n}{2} \rfloor} \beta_{n-1,i} + \sum_{i=0}^{\lfloor \frac{n}{2} \rfloor} \beta_{n-1,i+1} - \sum_{i=0}^{\lfloor \frac{n}{2} \rfloor} r_{n-1,i-1}. \quad (7)$$

Remark that

$$\tau_3(n-1) = \sum_{i=0}^{\lfloor \frac{n-1}{2} \rfloor} \beta_{n-1,i}.$$

The proof is an immediate consequence of the following considerations:

- the first summand in the right hand side of Identity (7) is equal to $\tau_3(n-1)$ if n is even. In case of n odd, instead, this term exceeds $\tau_3(n-1)$ by the Catalan number $C_{\frac{n-1}{2}}$,
- the second summand in the right hand side of Identity (7) is exactly $\tau_3(n-1)$,
- the third summand in the right hand side of Identity (7) exceeds the integer $\tau_3(n-1)$ exactly by $\beta_{n-1,0}$,
- the fourth summand in the right hand side of Identity (7) is obviously equal to $R(n-1)$.

□

Remark that an explicit evaluation of the integers $Q(n)$ and $R(n)$ can be obtained via the well known Hook Length Formula.

Previous considerations lead to the following result concerning the asymptotic behaviour of the sequence $\tau_3(n)$:

Proposition 8 *For every integer $n \geq 1$, we have:*

$$\frac{\tau_3(n)}{\tau_3(n-1)} < 3. \quad (8)$$

Moreover,

$$\lim_{n \rightarrow \infty} \frac{\tau_3(n)}{\tau_3(n-1)} = 3. \quad (9)$$

Proof Inequality (8) follows directly by previous considerations. In order to prove (9) we remark that, by Theorem (7), the ratio $\frac{\tau(n)}{\tau(n-1)}$ can be written as:

$$\frac{\tau_3(n)}{\tau_3(n-1)} = 3 - (U_1(n-1) + U_2(n-1) + U_3(n-1)),$$

where

$$\begin{aligned} U_1(n-1) &= \frac{E(n+1) \cdot C_{\frac{n-1}{2}}}{\tau_3(n-1)} \\ U_2(n-1) &= \frac{\beta_{n-1,0}}{\tau_3(n-1)} \\ U_3(n-1) &= \frac{R(n-1)}{\tau_3(n-1)}. \end{aligned}$$

The statement will be proved as soon as we show that each summand $U_i(n-1)$ goes to 0 as n goes to infinity. The numerator of each $U_i(n-1)$ is a sum of a suitable finite number f_λ . For every f_λ appearing in one of these expression, we can single out an appropriate subset S_λ of summands whose cardinality grows as n goes to infinity, such that:

$$\lim_{n \rightarrow \infty} \frac{f_\lambda}{\tau_3(n-1)} \leq \lim_{n \rightarrow \infty} \frac{f_\lambda}{\sum_{\mu \in S_\lambda} f_\mu} = 0.$$

For instance, consider the rectangular shape ρ over $n-1$ blocks, namely, the shape corresponding to the values $i = 0, k = \frac{n-1}{3}$. Obviously, such a shape exists only if $n-1 \equiv 0 \pmod{3}$. In this case, the family S_ρ consists of the shapes over $n-1$ cells and corresponding to the values $i = h, k = \frac{n-1}{3} - h$, where h ranges from 1 to $k = \frac{n-1}{6}$. By Hook Length Formula, we get:

$$\lim_{n \rightarrow \infty} \frac{f_\rho}{\tau_3(n-1)} \leq \lim_{n \rightarrow \infty} \frac{f_\rho}{\sum_{\mu \in S_\rho} f_\mu} = \lim_{n \rightarrow \infty} \frac{f_\rho}{\sum_{h=1}^{\lfloor \frac{n-1}{6} \rfloor} (h+1)^3 f_\rho} \leq \lim_{n \rightarrow \infty} \frac{1}{8^{\lfloor \frac{n-1}{6} \rfloor}} = 0.$$

The sets S_ν corresponding to the other summand of each of the numerators of the $U_i(n-1)$ can be described analogously. $\square$

As proved in [12], $\tau_3(n) = M_n$, where M_n is the n -th Motzkin number. Hence, these last arguments yield a combinatorial proof of Propositions 4 and 5 in [2].

4 The general case

In this section we extend the previous argumentations to the general case of standard Young tableaux with at most s columns. Define a matrix $\Gamma^{(s)} = (\gamma_{n,i}^{(s)})$ such that the (n,i) -th entry is the total number of standard Young tableaux with at most s columns and such that the difference between the length of the second and of the third column is i .

Also in this case, when $s \geq 4$ and $n \geq s$, the entries of this matrix satisfy a three-term recurrence. In fact, the following proposition can be proved by the same arguments used to prove Proposition 6:

Proposition 9 *The integers $\gamma_{n,0}^{(s)}$ satisfy the identity:*

$$\gamma_{n,0}^{(s)} = (s-2)\gamma_{n-1,0}^{(s)} + \gamma_{n-1,1}^{(s)} - \sum_{j=3}^{s-1} r_j^{(s)}(n-1,0). \quad (10)$$

Moreover, for every $1 \leq i \leq \lfloor \frac{n}{2} \rfloor$, we have:

$$\begin{aligned} \gamma_{n,i}^{(s)} &= \gamma_{n-1,i-1}^{(s)} + (s-2)\gamma_{n-1,i}^{(s)} + \gamma_{n-1,i+1}^{(s)} - r_1^{(s)}(n-1,i-1) \\ &\quad - \sum_{j=3}^{s-1} r_j^{(s)}(n-1,i), \end{aligned} \quad (11)$$

where $r_j^{(s)}(n-1,i)$ is the number of standard Young tableaux with at most s columns such that the difference between the length of the second and of the third column is i and the j -th and the $(j+1)$ -th columns have the same length.

$\square$

Similarly, exploiting the same arguments as in Proposition 8, we get:

Proposition 10 *The numbers $\tau_s(n)$ satisfy the recurrence:*

$$\tau_s(n) = s\tau_s(n-1) - E(n-1)C_{\frac{n-1}{2}} - \gamma_{n-1,0}^{(s)} - R^{(s)}(n-1), \quad (12)$$

where $R^{(s)}(n-1)$ is the sum of all the correction terms appearing in Formula (11).

□

Now we can easily deduce from these identities some information concerning the asymptotic behaviour of the sequence $\tau_s(n)$:

Proposition 11 *The sequence $\tau_s(n)$ satisfies the following properties:*

$$\frac{\tau_s(n)}{\tau_s(n-1)} < s$$

$$\lim_{n \rightarrow \infty} \frac{\tau_s(n)}{\tau_s(n-1)} = s.$$

□

Acknowledgments

We thank Elisa Pergola and Renzo Pinzani for many useful conversations.

References

- [1] M.Aigner, Catalan and other numbers: a recurrent theme, in *Algebraic Combinatorics and Theoretical Computer Science*, H.Crapo and D.Senato eds., Springer-Verlag, (2001), 347–390.
- [2] M.Aigner, Motzkin numbers, *Europ. J. Combinatorics* **19**, (1998), 663–675.
- [3] F.Bergeron, L.Favreau, D.Krob, Conjectures on the enumeration of tableaux of bounded height, *Discrete Math.* **139**, (1995), 463–468.

- [4] F.Bergeron, F.Gascon, Counting Young tableaux of bounded height, *J. Integer Seq.* **3**, No.1, Art. 00.1.7, 7 p., electronic only (2000)
- [5] J.S.Frame, G. De B.Robinson and R.Thrall, The hook graphs of the symmetric group, *Can. J. Math.*, **6**, (1954), 316–324.
- [6] G.Frobenius, Uber die charaktere de symmetrischen gruppen, *Preuss. Akad. Wiss. Sitz.*, (1900), 516–534.
- [7] I.Gessel, Symmetric functions and P-recursiveness, *Jour. of Comb. Th.*, Series A, **53**, (1990), 257–285.
- [8] D.Gouyou-Beauchamps, Standard Young tableaux of height 4 and 5, *Europ. J. Combinatorics.* **10**, (1989), 69–82.
- [9] D.Gouyou-Beauchamps, Codages par des mots et des chemins: problèmes combinatoires et algorithmiques, Ph.D. Thesis, University of Bordeaux I, 1985.
- [10] D.E.Knuth, “The art of computer programming: sorting and searching”, Vol. 3, Addison-Wesley (1998).
- [11] T.Nakayama, On some modular properties of irreducible representations of a symmetric group, I, II, *Jap. J. Math.*, **17**, (1940), 411–423.
- [12] A.Regev, Asymptotic values for degrees associated with strips of Young tableau, *Adv. in Math.*, **41**, (1981), 115–136.
- [13] N.J.A. Sloane, Encyclopedia of integer sequences, <http://www.research.att.com/~njas/sequences/>
- [14] R.P.Stanley, Differentiability finite power series, *Europ. J. Comb.*, **1**, (1980), 175–188.
- [15] A.Young, On quantitative substitutional analysis II, *Proc. Lond. Math. Soc.*, **34**, (1902), 361–397.